



INFECTIOUS DISEASE AND INFECTION PREVENTION AND CONTROL COVID-19 **COMMITTEE**

October 8, 2021

Bulletin #174b: ID-IPC Personal Protective Equipment (PPE) required when caring for **Suspect/Confirmed COVID-19 Patients**

COVID-19 Infection Prevention & Control (IPC) Guidance Documents

Personal Protective Equipment (PPE) required when caring for **Suspect/Confirmed COVID-19 Patients**

Prior to every patient interaction, healthcare workers (HCWs) have a responsibility to perform a Point of Care Risk Assessment (PCRA) to assess the infectious risk posed to themselves and others. A PCRA will help determine the correct PPE required to protect the HCW in their interaction with the patient and patient environment.

The use of Droplet/Contact Precautions (medical grade face mask, gown, gloves, eye protection) are recommended for all HCWs when caring for patients with suspect or confirmed COVID-19.

Full Precautions are followed when caring for suspect or confirmed COVID-19 patients and aerosolgenerating medical procedures (AGMPs) are being performed. A fit-tested and seal-checked N95 respirator is worn in place of a medical grade face mask in addition to gloves, gown, and eye protection.

An AGMP is a medical procedure that can generate aerosols as a result of artificial manipulation of a person's airway. The risk of transmission of infection may increase during AGMPs due to the potential to generate a high volume of respiratory aerosols that may be propelled over a longer distance than that involved in natural dispersion patterns.

To prevent transmission of infection associated with aerosols produced by AGMPs such as endotracheal intubation/extubation, CPR, etc. the Infection Prevention and Control Guidance for Performing an Aerosol Generating Medical Procedure During Patient Resuscitation and Other Scenarios guides the management of all patients receiving an AGMP in all patient care areas.

N95 respirators are required when providing care to suspect or confirmed COVID-19 patients following Full Precautions in the following settings:

- When providing direct patient care to a patient with suspect or confirmed COVID-19 during an AGMP.
- When in a patient care area/room where a suspect or confirmed COVID-19 patient is being managed and an AGMP is being performed.

















Please refer to the Infection Prevention & Control Guidance: Management of Suspect/Confirmed COVID-19 Patients posted under the IPC Resources COVID-19 Skyline Page.









